

WRIGHT-FISHER DIFFUSION WITH NEGATIVE MUTATION RATES

SOUMIK PAL

ABSTRACT. We study a family of n -dimensional diffusions taking values in the unit simplex of vectors with non-negative coordinates that add up to one. These processes satisfy stochastic differential equations which are similar to the ones for the classical Wright-Fisher diffusions, except that the “mutation rates” are now nonpositive. This model, suggested by Aldous, appears in the study of a conjectured diffusion limit for a Markov chain on Cladograms. The striking feature of these models is that the boundary is not reflecting, and we kill the process once it hits the boundary. We derive the explicit exit distribution from the simplex, and probabilistic bounds on the exit time. We also prove that these processes can be viewed as a “stochastic time-reversal” of a Wright-Fisher process of increasing dimensions and conditioned at a random time. A key idea in our proofs is a skew-product construction using certain one-dimensional diffusions called Bessel-square processes of negative dimensions which have been recently introduced by Göing-Jaeschke and Yor.

1. INTRODUCTION

An n -leaf Cladogram is an unrooted tree with $n \geq 4$ labeled leaves (vertices with degree one) and $(n - 2)$ other unlabeled vertices (internal branchpoints) of degree three (see below for a figure). The number of edges in such a tree is exactly $2n - 3$. Sometimes they are also referred to as phylogenetic trees. Aldous in [3] proposes the following model of a reversible Markov chain on the space of all n -leaf Cladograms which consists of removing a random leaf (and its incident edge) and reattaching it to one of the remaining random edges.

For a precise description we first define two operations on Cladograms. More details, with figures, can be found in [3].

- (i) To *remove a leaf* i . The leaf i is attached by an edge e_1 to a branchpoint b where two other edges e_2 and e_3 are incident. Delete edge e_1 and branchpoint b , and then merge the two remaining edges e_2 and e_3 into a single edge e . The resulting tree has $2n - 5$ edges.
- (ii) To *add a leaf* to an edge f . Create a branchpoint b' which splits the edge f into two edges f_2, f_3 and attach the leaf i to branchpoint b' via a new edge f_1 . This restores the number of leaves and edges to the tree.

Let $\mathbf{T}_n$ denote the finite collection of all n -leaf Cladograms. Write $\mathbf{t}' \sim \mathbf{t}$ if $\mathbf{t}' \neq \mathbf{t}$ and $\mathbf{t}'$ can be obtained from $\mathbf{t}$ by following the two operations above for some choice of i and f . Thus a $\mathbf{T}_n$ valued chain can be described by saying:

Date: January 31, 2012.

2000 Mathematics Subject Classification. 60G99; 60J05; 60J60; 60J80.

Key words and phrases. Wright-Fisher diffusion, Markov chain on Cladograms, Continuum Random Tree, Bessel processes of negative dimension.




FIGURE 1. A 7-leaf Cladogram

remove leaf i uniformly at random, then pick edge f at random and re-attach i to f . If we assume every edge to be of unit length, then it also involves resizing the edge length after every operation. In particular the transition matrix of this Markov chain is

$$P(\mathbf{t}, \mathbf{t}') = \begin{cases} \frac{1}{n(2n-5)}, & \text{if } \mathbf{t}' \sim \mathbf{t} \\ \frac{n}{n(2n-5)}, & \text{if } \mathbf{t}' = \mathbf{t}. \end{cases}$$

This leads to a symmetric, aperiodic, and irreducible finite state space Markov chain. Schweinsberg [16] proved that the relaxation time for this chain is $O(n^2)$, improving a previous result in [3].

In his webpage [2] Aldous asks the following question: what is an appropriate diffusion limit of this Markov chain? The invariant distribution for the Markov chain on n -leaf Cladograms is clearly the Uniform distribution. It is known, see Aldous [1], that the sequence of Uniform distributions on n -leaf Cladograms converge weakly to the law of the (Brownian) Continuum Random Tree (CRT). Hence, it is natural to look for an appropriate Markov process on the support of the CRT which can be thought of as a limit of the sequence of Markov chains described above. At this point it is important to understand that the support of the CRT consists of compact real trees with a measure describing the distribution of leaves. These trees are called continuum trees. For a formal definition of these concepts we refer the reader to the seminal work by Aldous in [1]. However, for an intuitive visualization, one should think of a typical continuum tree as a compact metric space on which branchpoints are dense and all edges are infinitesimally small. This implies that the Markov process that mimics the operation of removing and inserting a new leaf on a continuum tree should not jump, in other words, we can call it a diffusion.

A detailed description of this diffusion on continuum trees is forthcoming in Pal [13]. In this article we consider several important features of this limiting diffusion that are of interest by themselves and provide bedrock for the followup construction.

Consider the branchpoint b in the 7-leaf Cladogram $\mathbf{t}$ in Figure 1. It divides the collection of leaves naturally into three sets. Let $X(\mathbf{t}) = (X_1, X_2, X_3)(\mathbf{t})$ denote the vector of proportion of leaves in each set. The corresponding number of edges in these sets are $(2nX_1 - 1, 2nX_2 - 1, 2nX_3 - 1)$. For example, at time zero in our given tree, going clockwise from the right we have $X(0) = (3/7, 2/7, 2/7)$.

Let $\mathbb{S}_n$ denote the unit simplex

$$(1) \quad \mathbb{S}_n = \left\{ x \in \mathbb{R}^n : x_i \geq 0 \text{ for all } i \text{ and } \sum_{i=1}^n x_i = 1 \right\}.$$

Some simple algebra will reveal that for any point $x = (x_1, x_2, x_3)$ in $\mathbb{S}_3$, given $X(\mathbf{t}) = x$, the difference $X_1(\mathbf{t}') - X_1(\mathbf{t})$ can only take values in $\{-1/n, 0, 1/n\}$ with corresponding probabilities

$$q_{x_1} = x_1 \frac{2n(1-x_1)-2}{2n-5}, \quad 1-p_{x_1}-q_{x_1}, \quad p_{x_1} = (1-x_1) \frac{2nx_1-1}{2n-5}.$$

Thus

$$\begin{aligned} E(X_1(\mathbf{t}') - X_1(\mathbf{t}) \mid X(\mathbf{t}) = x) &= \frac{1}{n} \frac{2x_1 - (1-x_1)}{2n-5} \approx -\frac{1}{n^2} \frac{1}{2} (1-3x_1) \\ (2) \quad E((X_1(\mathbf{t}') - X_1(\mathbf{t}))^2 \mid X(\mathbf{t}) = x) &= \frac{1}{n^2} \frac{4nx_1(1-x_1) - x_1 - 1}{2n-5} \\ &\approx \frac{1}{n^2} 2x_1(1-x_1). \end{aligned}$$

If we take scaled limits, as n goes to infinity, of the first two conditional moments (the mixed moments can be similarly verified), it is intuitive (and follows by standard tools) that as n goes to infinity, this Markov chain (run at $n^2/2$ speed) will converge to a diffusion with a generator

$$(3) \quad \frac{1}{2} \sum_{i,j=1}^3 x_i (1\{i=j\} - x_j) \frac{\partial^2}{\partial x_i \partial x_j} - \frac{1}{2} \sum_{i=1}^n \frac{1}{2} (1-3x_i) \frac{\partial}{\partial x_i}.$$

The generator written as above is similar to the generator for the well-known diffusion limit of the Wright-Fisher (WF) Markov chain models in population genetics. The WF model is one of the most popular models in population genetics. This is a multidimensional Markov chain which keeps track of the vector of proportions of certain genetic traits in a population of non-overlapping generations. A good source for an introduction to these models is Chapter 1 in the book by Durrett [6]. For computational purposes one often takes recourse to a diffusion approximation, which, in its standard form, leads to a family of diffusions parametrized by n “mutation rates”. The state space of the diffusion is given by $\mathbb{S}_n$ and is parametrized by a vector $(\delta_1, \dots, \delta_n)$ of nonnegative entries. A weak solution of the WF diffusion with parameters $\delta = (\delta_1, \dots, \delta_n)$ solves the following stochastic differential equation for $i = 1, 2, \dots, n$:

$$(4) \quad dJ_i(t) = \frac{1}{2} (\delta_i - \delta_0 J_i(t)) dt + \sum_{j=1}^n \tilde{\sigma}_{i,j}(J) d\beta_j(t), \quad \delta_0 = \sum_{i=1}^n \delta_i.$$

Here $\beta = (\beta_1, \dots, \beta_n)$ is a standard multidimensional Brownian motion and the diffusion matrix $\tilde{\sigma}$ is given by

$$(5) \quad \tilde{\sigma}_{i,j}(x) = \sqrt{x_i} (1\{i=j\} - \sqrt{x_i x_j}), \quad 1 \leq i, j \leq n.$$

We define the Wright-Fisher diffusion with *negative mutation rates* to be a family of n -dimensional diffusions, parametrized by n nonnegative parameters $\delta = (\delta_1, \dots, \delta_n)$, which is a weak solution of the following differential equation:

$$(6) \quad d\mu_i(t) = -\frac{1}{2} (\delta_i - \delta_0 \mu_i(t)) dt + \sum_{j=1}^n \tilde{\sigma}_{i,j}(\mu) d\beta_j(t), \quad \delta_0 = \sum_{i=1}^n \delta_i.$$

The initial condition $\mu(0)$ is in the interior of $\mathbb{S}_n$ and the process has a drift that pushes it outside the simplex. We will show later that the process is sure to hit the boundary of the simplex at which point we stop it. In the next section we will

explicitly construct a weak solution of (6). The uniqueness in law of such a solution until it hits the boundary follows since the drift and the diffusion coefficients are smooth (hence, Lipschitz) inside the open unit simplex. The law of this process will then be denoted uniquely by $\text{NWF}(\delta_1, \dots, \delta_n)$.

Equivalently this process can be identified by its Markov generator. Expanding $\tilde{\sigma}\tilde{\sigma}'$ and using the fact that $\sum_{i=1}^n x_i = 1$, we get

$$(7) \quad \mathcal{A}_n = \frac{1}{2} \sum_{i,j=1}^n x_i (1\{i=j\} - x_j) \frac{\partial^2}{\partial x_i \partial x_j} - \sum_{i=1}^n \frac{1}{2} (\delta_i - \delta_0 x_i) \frac{\partial}{\partial x_i}.$$

which identifies (3) as the generator for $\text{NWF}(1/2, 1/2, 1/2)$.

In this text we focus on properties of NWF models as a family of diffusions on the unit simplex and explore some of their properties that are important in the context of the Markov chain model on Cladograms.

Part (1). We show that, just like Wright-Fisher diffusions (see [12]), the NWF processes can be recovered from a far simpler class of models, the Bessel-Square (BESQ) processes with negative dimensions. A comprehensive treatment of BESQ processes can be found in the book by Revuz & Yor [15]. This family of one dimensional diffusions is indexed by a single real parameter θ (called the dimension) and are solutions of the stochastic differential equations

$$(8) \quad Z(t) = x + 2 \int_0^t \sqrt{|Z(s)|} d\beta(s) + \theta t, \quad x \geq 0, \quad t \geq 0,$$

where β is a one dimensional standard Brownian motion. We denote the law of this process by Q_x^θ . It can be shown that the above SDE admits a unique strong solution until it hits the origin. The classical model only admits parameter θ to be non-negative. However, an extension, introduced by Göing-Jaeschke & Yor [7], allows the parameter θ to be negative. It is important to note that Q_x^θ is the diffusion limit of a Galton-Watson branching process with a $|\theta|$ rate of immigration (for $\theta \geq 0$) or emigration (for $\theta < 0$).

In Section 3 we show that the $\text{NWF}(\delta_1, \dots, \delta_n)$ law, starting at $(x_1, \dots, x_n)$, can be recovered via a stochastic time-change from a collection of n independent processes with laws $Q_{x_i}^{-2\delta_i}$, $i = 1, \dots, n$, and dividing each coordinate by the total sum. For the corresponding discrete models this is usually referred to as Poissonization.

In this article we utilize this relationship to infer several properties about the NWF processes. For example, we prove that these diffusions, almost surely, hit the boundary of the simplex. We derive the explicit exit density supported on the union of the boundary walls in Theorem 9.

Part (2). We also prove an interesting duality relationship between WF and NWF models. To describe the duality relationship we let the NWF continue in the lower dimensional simplex when any of the coordinates hit zero. Thus, every time a coordinate hits zero, the dimension of the process gets reduced by one, and ultimately the process is absorbed at the scalar one. Such a process can be obtained by running a WF model with appropriate parameters that initially starts with dimension one and value 1. At independent random times, the dimension of the process increases by one, and the newly added coordinate is initialized at zero. Finally we condition on the values of the process at a chosen random time.

The resulting process, backwards in time and suitably time-changed, is the original NWF model.

Part (3). The time that the NWF process takes to exit the simplex is a crucial quantity due to a reason which we describe below. We keep our exposition mostly verbal without going into too much detail since the details require considerable formalism from the theory of continuum trees and will be discussed elsewhere. In [13] we show how **Part (1)** points towards a Poissonization of the entire Aldous Markov chain which is simpler for considering scaled limits. The Poissonized version of the Markov chain on n -leaf Cladograms stipulates: every existing leaf has an Exponential clock of rate 2 attached to it which determines the instances of their deaths, and every existing edge has an independent Exponential clock of rate 1 attached to it at which point the edge is split and a new pair of vertices (one of which is a leaf) is introduced. It is an easy verification that the rates are consistent with the BESQ limit that we claimed in **Part (1)** above. Hence, one would expect that the limit of the Poissonized chains on continuum trees, normalized to give a leaf-mass measure one, and suitably time-changed would give the conjectured Aldous diffusion. This is the strategy followed in [13].

Now, the Poissonized chain has some beautiful and interesting structure. Please see [1] for the details about continuum trees that we use below. A continuum tree $\mathbb{T}$ comes with its associated (infinite) length measure (analogous to the Lebesgue measure) and a leaf-mass probability measure which describes how the leaves are distributed on it. We will denote the length measure by $\mathbf{Leb}(\mathbb{T})$ and the leaf-mass probability measure by $\mu(\mathbb{T})$. Suppose we sample n iid elements from $\mu(\mathbb{T})$ and draw the tree generated by them, it produces an n -leaf Cladogram with edge-lengths (or, a proper n -tree, according to [1]). Thus, by using the fact that the continuum tree is compact, one can approximate a continuum tree by a sequence of n -leaf Cladograms.

Now consider an n -leaf Cladogram for a very large n , and further consider m internal branchpoints. For example, in Figure 1, we have three branchpoints $\{a, b, c\}$ in a 7-leaf Cladogram. These branchpoints generate a *skeleton* subtree of the original tree and partition the leaves as *internal* or *external* to the skeleton. The components of the vector of external leaf masses grow as independent continuous time, binary branching, Galton-Watson branching processes with a rate of branching / dying 2 and a rate of emigration 1. Note that this is consistent with the diffusion limit as BESQ with $\theta = -1$. As the Markov chain (Poissonized or not) proceeds, there comes a time when one of these external leaf masses gets exhausted. When this happens, one of the internal branchpoints becomes a leaf. The distribution of every coordinate of external leaf-masses at this exit time is derived in **Part (2)**. Until this time, supported on the skeleton, new subtrees can grow and decay. We show in [13], the dynamics of the sizes of these subtrees on the internal part can be modeled as the age process of a chronological splitting tree. Chronological splitting trees are special kind of biological trees where an individual lives up to a certain (possibly non-Exponential) lifetime and produces children at rate one during that lifetime. Her children behave in an identical manner with an independent and identically distributed lifetime of their own. The age process refers to the point process of current ages of the existing members in the family. More details about splitting trees can be found in the article by Lambert [10].

When one of the internal vertices gets *exposed*, the above dynamics breaks down and we need to find a slightly different set of internal vertices to proceed. Hence, it is important to derive estimates of the times at which this change happens. We provide quantitative bounds on the value of this stopping time under the special situation of symmetric choice of parameters which is the case at hand.

The article is divided as follows. Our main tool in this analysis is to establish a relationship between NWF processes and Bessel-square processes of negative dimensions, much in the spirit of Pal [12]. This has been done in Section 3 where we also establish Theorem 7. The relevant results about BESQ processes have been listed in Section 2. Most of these results are known and appropriate citations have been provided. Proofs of the rest have been done in the Appendix. Exact computations of exit density from the simplex has been done in Section 4. Estimates of the exit time have been established in Section 5.

2. SOME RESULTS ABOUT BESQ PROCESSES

The Bessel-Square processes of negative dimensions $-\theta$, where $\theta \geq 0$, are one dimensional diffusions which are the unique strong solution of the SDE

$$(9) \quad X(t) = x - \theta t + 2 \int_0^t \sqrt{X(s)} d\beta(s), \quad t \leq T_0,$$

where T_0 is the first hitting time of zero for the process X and x is a positive constant. The process is absorbed at zero. We will denote the law of this process $Q_x^{-\theta}$ just as BESQ of a positive dimension θ will be denoted by Q_x^θ .

The following collection of results is important for us. All the proofs can be found in the article by Göing-Jaesche and Yor [7].

Lemma 1 (Time-reversal). *For any $\theta > -2$ and any $x > 0$, $Q_x^{-\theta}(T_0 < \infty) = 1$, while, for $\theta \geq 2$, one has $Q_x^\theta(T_0 < \infty) = 0$.*

Moreover the following equality holds in distribution:

$$(10) \quad (X(T_0 - u), u \leq T_0) = (Y(u), u \leq L_x),$$

where Y has law $Q_0^{4+\theta}$ and L_x is the last hitting time of x for the process Y .

In particular,

- (i) both L_x and T_0 are distributed as $x/2G$, where G is a Gamma random variable with parameter $(\theta/2 + 1)$.
- (ii) The transition probabilities $p_t^\theta(x, y)$ for $x, y > 0$ satisfy the identity

$$p_t^{-\theta}(x, y) = p_t^{4+\theta}(y, x).$$

The following results have been proved in the Appendix.

Lemma 2. *The scale function for $Q^{-\theta}$, $\theta \geq 0$, is given by the function*

$$s(x) = x^{\theta/2+1}, \quad x \geq 0.$$

Moreover,

- (i) *The origin is an exit boundary for the diffusion and not an entry.*
- (ii) *The change of measure*

$$x^{-\theta/2-1} Q_x^{-\theta} \left(X(t)^{\theta/2+1} 1(\cdot) \right)$$

on the σ -algebra generated by the process up to time t is consistent for various t and is the law of $Q_x^{4+\theta}$. Thus, we say $Q_x^{4+\theta}$ is $Q_x^{-\theta}$ conditioned never to hit zero.

The previous fact is the generalization of the well-known observation that Brownian motion, conditioned never to hit the origin, has the law of the three-dimensional Bessel process.

Lemma 3. *Let $\{Z(t), t \geq 0\}$ denote a BESQ process of dimension θ for some $\theta > 2$. Then*

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\log(1/\epsilon)} \int_{\epsilon}^t \frac{du}{Z(u)} = \frac{1}{\theta - 2}, \quad \text{for all } t > 0.$$

3. CHANGING AND REVERSING TIME

Our objective in this section is to establish a time-reversal relationship between NWF and WF models.

Theorem 4. *Let $z_1, \dots, z_n$ and $\theta_1, \dots, \theta_n$ be nonnegative constants. Let $Z = (Z_1, \dots, Z_n)$ be a vector of n independent BESQ processes of dimensions $-\theta_1, \dots, -\theta_n$ respectively, starting from $(z_1, \dots, z_n)$. Let ζ be the sum $\sum_{i=1}^n Z_i$.*

Define

$$T_i = \inf \{t \geq 0 : Z_i(t) = 0\}, \quad \tau = \wedge_{i=1}^n T_i.$$

Then, there is an n -dimensional diffusion μ , satisfying the SDE in (6) for $NWF(\theta_1/2, \dots, \theta_n/2)$, for which the following equality holds:

$$(11) \quad Z_i(t \wedge \tau) = \zeta(t \wedge \tau) \mu_i(4C_t), \quad 1 \leq i \leq n, \quad C_t = \int_0^{t \wedge \tau} \frac{ds}{\zeta(s)}.$$

Thus, in particular, equation (6) admits a weak solution for all non-negative parameters $(\delta_1, \dots, \delta_n)$.

Proof. The proof is almost identical to the case of WF model as shown in [12, Proposition 11] with obvious modifications. For example, unlike the WF case, the time-change clock is no longer independent of the NWF process. We outline the basic steps below.

We know from (9) that

$$dZ_i(t \wedge \tau) = -\theta_i d(t \wedge \tau) + 2\sqrt{Z_i} d\beta_i(t \wedge \tau), \quad i = 1, 2, \dots, n.$$

Define $\theta_0 = \sum_{i=1}^n \theta_i$. Let $V_i(t) = Z_i/\zeta(t)$ for $t \leq \tau$. Then by Itô's rule we get:

$$(12) \quad dV_i(t \wedge \tau) = -\zeta^{-1} [\theta_i - \theta_0 V_i] d(t \wedge \tau) + \sqrt{V_i(1 - V_i)} dM_i(t),$$

where

$$(13) \quad dM_i(t) = \frac{2\zeta^{-1/2}}{\sqrt{1 - V_i}} \sum_{j=1}^n \left(1\{i = j\} - \sqrt{V_i V_j}\right) d\beta_j(t \wedge \tau),$$

and $\langle M_i \rangle(t) = 4C_t$.

Let $\{\rho_u, u \geq 0\}$ be the inverse of the increasing function $4C_t$. Applying this time-change to the SDE for V_i in (12), we get

$$(14) \quad d\mu_i(t) = -\frac{1}{4} [\theta_i - \theta_0 \mu_i] dt + \sqrt{\mu_i(1 - \mu_i)} \widetilde{W}_i(t),$$

where $\widetilde{W}_i$ is the Dambis-Dubins-Schwarz (DDS; see [15, p. 181]) Brownian motion associated with M_i . This turns out to be the SDE for $NWF(\theta_1/2, \dots, \theta_n/2)$. $\square$

Let $\theta_1, \theta_2, \dots, \theta_n$ be nonnegative and $z_1, z_2, \dots, z_n$ be positive constants. For $i = 1, 2, \dots, n$ define independent random variables $(G_1, \dots, G_n)$ where G_i is distributed as $\text{Gamma}(\theta_i/2 + 1)$. Let

$$(15) \quad R_i = \frac{z_i}{2G_i}, \quad i = 1, 2, \dots, n.$$

Also, independent of $(G_1, \dots, G_n)$, let $Y_1, Y_2, \dots, Y_n$ be n independent BESQ processes of positive dimensions $(4 + \theta_1), (4 + \theta_2), \dots, (4 + \theta_n)$ respectively, all of which are starting from zero.

For any permutation π of n labels, condition on the event

$$(16) \quad R_{\pi_1} > R_{\pi_2} > \dots > R_{\pi_n}, \quad \text{and let } R^* = R_{\pi_2}.$$

We now construct the following n dimensional process $(X_1, \dots, X_n)$:

$$(17) \quad X_i(t) = Y_i((t - R^* + R_i)^+), \quad t \geq 0.$$

Notice that at time $t = 0$, every X_i is at zero except the π_1 th.

Let $S(t)$ denote the total sum process $\sum_{i=1}^n X_i(t)$. Note that $S(t) > 0$ for all $t \geq 0$ with probability one. Define the process

$$(18) \quad C_t := \int_0^t \frac{du}{S(u)}, \quad t > 0.$$

The process C_t is finite almost surely for every t (unfortunately, we cannot define $R^* = R_{\pi_1}$ precisely because C_t will be infinity; see Lemma 3). Let A denote the inverse function of the continuous increasing function $4C$. That is

$$(19) \quad A_t = \inf \{u \geq 0 : 4C_u \geq t\}, \quad t \geq 0.$$

Lemma 5. *There is an n -dimensional diffusion ν such that the following time-change relationship holds*

$$(20) \quad \nu_i(t) = \frac{X_i}{S}(A_t), \quad \text{or} \quad X_i(t) = S(t)\nu_i(4C_t) \quad t \geq 0.$$

The distribution of ν is supported on the unit simplex

$$\mathbb{S}_n = \{x_i \geq 0 : x_1 + x_2 + \dots + x_n = 1\}.$$

Conditional on the values of $G_1, \dots, G_n$ and the process S , the law of ν can be described as below.

Let π be any permutation of n labels. On the event $R_{\pi_1} > R^ = R_{\pi_2} > \dots > R_{\pi_n}$. Let $V_2 < \dots < V_n$ be defined by*

$$A_{V_i} = R^* - R_{\pi_i}, \quad \text{or, equivalently} \quad 4C_{R^* - R_{\pi_i}} = V_i.$$

Note that $V_2 = 0$.

For $i \geq 2$ and $V_i \leq t \leq V_{i+1}$, the process ν is zero on all coordinates except $(\pi_1, \dots, \pi_i)$. The process $\nu(\pi_1, \dots, \pi_i)$, given the history of the process till time V_i (and the G_i 's and S) is distributed as the classical Wright-Fisher diffusion starting from

$$\frac{1}{S}(X_{\pi_1}, \dots, X_{\pi_i})(A_{V_i}) = \frac{1}{S}(X_{\pi_1}, \dots, X_{\pi_i})(R^* - R_{\pi_i}),$$

and with parameters $(\gamma_{\pi_1}, \dots, \gamma_{\pi_i})$ where

$$\gamma_j = \theta_j/2 + 2, \quad j = 1, 2, \dots, n.$$

Proof. The Gamma random variables $G_1, \dots, G_n$ are independent of the BESQ process $Y_1, \dots, Y_n$. Thus, conditional on $G_1, \dots, G_n$, the vector of processes $(X_1, \dots, X_n)$ has the following description. For

$$R^* - R_{\pi_i} \leq t \leq R^* - R_{\pi_{i+1}}, \quad i \geq 2,$$

all coordinates other than the π_1 th, π_2 th, $\dots$, π_i th are zero. And, $(X_{\pi_1}, \dots, X_{\pi_i})$, conditioned on the past, are independent BESQ processes of dimensions $(4 + \theta_{\pi_1}, \dots, 4 + \theta_{\pi_i})$ and starting from $(X_{\pi_1}, \dots, X_{\pi_i})(R^* - R_{\pi_i})$.

Thus, on this interval of time, the existence of the process ν , identifying its law as the WF law, and the claimed independence from the process S , all follow from [12, Prop. 11]. The proof of the Lemma now follows by combining the argument over the distinct intervals. $\square$

Lemma 6. *Consider the set-up in (15), (17), and (19). Let $Z_1, Z_2, \dots, Z_n$ be n stochastic processes defined such that $\{Z_i(t), 0 \leq t \leq R^*\}$ is the time-reversal of the process $\{X_i(t), 0 \leq t \leq R^*\}$, conditioned on $X_i(R^*) = z_i$. That is, conditioned on $X_i(R^*) = z_i$ for every i ,*

$$Z_i(t) = X_i(R^* - t) = Y_i(R_i - t)^+, \quad \text{for } 0 \leq t \leq R^*.$$

Then $(Z_1, \dots, Z_n)$ are independent BESQ processes of dimensions $-\theta_1, \dots, -\theta_n$, starting from $z_1, \dots, z_n$, and absorbed at the origin.

Proof. It suffices to prove the following:

Claim. Let $\{Y(t), t \geq 0\}$ denote a BESQ process of dimension $(4 + \theta)$ starting from 0. Fix a $z > 0$. Let T be distributed as $z/2G$, where G is a Gamma random variable with parameter $(\theta/2 + 1)$. Then, conditioned on $T = l$ and $Y(l) = z$, the time-reversed process $\{Y((l - s)^+), 0 \leq s < \infty\}$ is distributed as $Q_z^{-\theta}$, absorbed at the origin, conditioned on $T_0 = l$. Here T_0 is the hitting time of the origin for $Q_z^{-\theta}$.

Once we prove this claim, the lemma follows since the law of T_0 is exactly $z/2G$. See Lemma 1.

Proof of Claim. For the case of $\theta = 0$, this is proved in [14, p. 447]. The general proof is exactly similar and we outline just the steps and give references within [14] for the details.

For any $\theta \in \mathbb{R}$, $t > 0$, $x, y \geq 0$, let $Q_{x \rightarrow y}^{\theta, t}$ denote the law of the BESQ bridge of dimension θ , length t , from points x to y . That is to say, if Y follows Q_x^θ , then $Q_{x \rightarrow y}^{\theta, t}$ is the law of the process $\{Y(s), 0 \leq s \leq t\}$ conditioned on the event $\{Y(t) = y\}$.

Now, BESQ bridges satisfies time-reversal [14, p. 446]. Thus, if we define $\hat{P}$ to be the P -distribution of a process $\{X(t - s), 0 \leq s \leq t\}$, then $Q_{x \rightarrow y}^{\theta, t} = \hat{Q}_{y \rightarrow x}^{\theta, t}$.

We consider the case when the dimension is $(4 + \theta)$, $\theta \geq 0$, $x = 0, y = z > 0$. Then

$$Q_{0 \rightarrow z}^{4+\theta, t} = \hat{Q}_{z \rightarrow 0}^{4+\theta, t}$$

Now, from Lemma 2 (also see [14, Section 3, p. 440]) we know that $Q_z^{4+\theta}$ is $Q_z^{-\theta}$ conditioned never to hit zero (or equivalently, $Q_z^{-\theta}$ can be interpreted as $Q_z^{4+\theta}$ conditioned to hit zero). Since the origin is an exit distribution for $Q_z^{-\theta}$ and not an entry (Lemma 2; see [14, p. 441] for the details of these definitions), the conditional law $Q_{z \rightarrow 0}^{4+\theta, t}$ is nothing but $Q_z^{-\theta}$, conditioned on $T_0 = t$. This completes the proof. $\square$

The following is a more precise statement.

Let $(z_1, \dots, z_n)$ be a point in the n -dimensional unit simplex $\mathbb{S}_n$. Fix n nonnegative parameters $\delta_1, \dots, \delta_n$. Let $G_1, \dots, G_n$ denote n independent Gamma random variables with parameters $\delta_1 + 1, \dots, \delta_n + 1$ respectively. Define $R_i = z_i/2G_i$.

For any permutation π of n labels, condition on the event $R_{\pi_1} > R_{\pi_2} > \dots > R_{\pi_n}$, and let $R^* = R_{\pi_2}$.

Define the continuous process S by prescribing $S(0) = Z_1(R_{\pi_1} - R^*)$ where Z_1 is distributed as $Q_0^{4+2\delta_{\pi_1}}$; and for any t such that

$$R^* - R_{\pi_i} \leq t \leq R^* - R_{\pi_{i+1}}, \quad i \geq 2, \quad R_{\pi_{n+1}} = 0.$$

given the history, the process is distributed as a Bessel-square process of dimension $\sum_{j=1}^i (4 + 2\delta_{\pi_j})$ starting from $S(R^* - R_{\pi_i})$.

Define the stochastic clocks

$$C_t = \int_0^t \frac{du}{S(u)}, \quad \widehat{C}_t = \int_{R^*-t}^{R^*} \frac{du}{S(u)}, \quad 0 \leq t \leq R^*.$$

and let $\widehat{A}_t$ denote the inverse function of $4\widehat{C}_t$. Let $V_2 < \dots < V_n$ be defined by $4C_{R^*-R_{\pi_i}} = V_i$. Note that $V_2 = 0$. The 4 is a standardization constant that appears due to the factor of 2 in the diffusion coefficient in (8).

Define an n -dimensional process ν given $R_1, \dots, R_n$ and the process S . For $i \geq 2$ and $V_i \leq t \leq V_{i+1}$, the process ν is zero on all coordinates except possibly at indices $(\pi_1, \dots, \pi_i)$. At time zero, the process starts at the vector that is 1 in the π_1 th coordinate and zero elsewhere.

Conditioned on the history till time V_i , the process $\{\nu(\pi_1, \dots, \pi_i)(t), V_i \leq t \leq V_{i+1}\}$ is distributed as the classical Wright-Fisher diffusion starting from $\nu(\pi_1, \dots, \pi_i)(V_i)$ and with parameters $(\gamma_{\pi_1}, \dots, \gamma_{\pi_i})$ where

$$\gamma_j = \delta_j + 2, \quad j = 1, 2, \dots, n.$$

Finally, consider the conditional law of the process, conditioned on the event

$$S(R^*)\nu_i(4C_{R^*}) = z_i, \quad \text{for all } i = 1, 2, \dots, n.$$

Theorem 7. *Define the time-reversed process*

$$\mu(t) = \nu\left(\widehat{A} \circ 4C_{R^*-t}\right),$$

where $\circ$ denotes composition. Then this conditional stochastic time-reversed process, until the first time any of the coordinates hit zero, has a marginal distribution (when G_i 's and S are integrated out) NWF($\delta_1, \dots, \delta_n$) starting from $(z_1, \dots, z_n)$.

Proof of Theorem 7. We start with given values of $R_{\pi_1} > R_{\pi_2} > \dots > R_{\pi_n}$ and the process S . Applying equation (20) in Lemma 5 to obtain the processes $(X_1, \dots, X_n)$ defined by

$$X_i(t) = S(t)\nu_i(4C_t), \quad 0 \leq t \leq R^*.$$

Then, the vector $(X_1, X_2, \dots, X_n)$ has the law prescribed by (17).

Now we apply Lemma 6 to obtain $(Z_1, \dots, Z_n)$ by conditioning $(X_1, \dots, X_n)$ and reversing time. Finally the construction in Theorem 4 gives us the vector $(\mu_1, \dots, \mu_n)$ from $(Z_1, \dots, Z_n)$ as desired. $\square$

4. EXIT DENSITY

Let $Z_1, Z_2, \dots, Z_n$ be independent BESQ processes of dimensions $-\theta_1, \dots, -\theta_n$, where each $\theta_i \geq 0$. We assume that at time zero, the vector $\mathbf{Z} = (Z_1, \dots, Z_n)$ starts from a point $\mathbf{z} = (z_1, \dots, z_n)$ where every $z_i > 0$. Define T_i to be the first hitting time of zero for the process Z_i , and let $\tau = \wedge_i T_i$ denotes the first time any coordinate hits zero. We would like to determine the joint distribution of $(\tau, \mathbf{Z}(\tau))$.

Note that since each T_i is a continuous random variable, the minimum is attained at a unique i . Thus, for a fixed $1 \leq i \leq n$, conditioned on the event $\tau = T_i$, the distribution of $Z_i(\tau)$ is the unit mass at zero, and the distribution of every other $Z_j(\tau)$ is supported on $(0, \infty)$. Now, let h_i denote the density of the stopping time T_i on $(0, \infty)$ and let $q_t^{-\theta}$ refers to the transition density of $Q^{-\theta}$. It follows that for any $a_j > 0$, $j \neq i$, we get

$$\begin{aligned} & P(\tau = T_i, \tau \leq t, Z_j(\tau) \geq a_j, \text{ for all } j \neq i) \\ &= P(T_i \leq t, T_j > T_i, Z_j(T_i) \geq a_j, \text{ for all } j \neq i) \\ &= \int_0^t h_i(s) \prod_{j \neq i} P(T_j > s, Z_j(s) \geq a_j) ds = \int_0^t h_i(s) \prod_{j \neq i} P(Z_j(s) \geq a_j) ds, \quad \text{since } a_j > 0. \\ &= \int_0^t h_i(s) \left[\prod_{j \neq i} \int_{a_j}^{\infty} q_s^{-\theta_j}(z_j, y_j) dy_j \right] ds. \end{aligned}$$

Our first job is to find closed form expressions of the integral above. To do this we start by noting that T_i is distributed as $z_i/2G_i$ (see Lemma 1), where G_i is a Gamma random variable with parameter $(4 + \theta_i)/2 - 1 = \theta_i/2 + 1$. That is, the density of G_i is supported on $(0, \infty)$ and is given by

$$\frac{y^{\theta_i/2}}{\Gamma(\theta_i/2 + 1)} e^{-y}.$$

It follows that

$$h_i(s) = \frac{(z_i/2)^{\theta_i/2+1}}{\Gamma(\theta_i/2 + 1)} s^{-\theta_i/2-2} e^{-z_i/2s}, \quad 0 \leq s < \infty.$$

On the other hand, it follows from time reversal (Lemma 1) that $q_s^{-\theta_j}(z_j, y_j) = q_s^{4+\theta_j}(y_j, z_j)$. For any positive a , the transition density $q_s^a(y, z)$ is explicitly known (see, e.g., in [12]) to be $s^{-1}f(z/s, a, y/s)$, where $f(\cdot, k, \lambda)$ is the density of a non-central Chi-square distribution with k -degrees of freedom and a noncentrality parameter value λ . In particular, it can be written as a Poisson mixture of central Chi-square (or, Gamma) densities. Thus, we have the following expansion

$$(21) \quad q_s^{-\theta_j}(z_j, y_j) = q_s^{4+\theta_j}(y_j, z_j) = s^{-1} \sum_{k=0}^{\infty} e^{-y_j/2s} \frac{(y_j/2s)^k}{k!} g_{\theta_j+4+2k}(z_j/s),$$

where g_r is the Gamma density with parameters $(r/2, 1/2)$. That is

$$g_r(x) = \frac{2^{-r/2} x^{r/2-1}}{\Gamma(r/2)} e^{-x/2}, \quad x \geq 0.$$

Now, define

$$\vec{y}_i = \sum_{j \neq i} y_j, \quad \vec{\theta}_i = \sum_{j \neq i} \theta_j, \quad \vec{z}_i = \sum_{j \neq i} z_j$$

Thus

$$\begin{aligned}
h_i(s) \prod_{j \neq i} q_s^{-\theta_j}(z_j, y_j) &= \frac{(z_i/2)^{\theta_i/2+1}}{\Gamma(\theta_i/2+1)} s^{-\theta_i/2-2} e^{-z_i/2s} \prod_{j \neq i} s^{-1} \sum_{k=0}^{\infty} e^{-y_j/2s} \frac{(y_j/2s)^k}{k!} g_{\theta_j+4+2k}(z_j/s) \\
&= \frac{(z_i/2)^{\theta_i/2+1}}{\Gamma(\theta_i/2+1)} s^{-\theta_i/2-2} e^{-z_i/2s} \prod_{j \neq i} s^{-1} \sum_{k=0}^{\infty} e^{-y_j/2s} \frac{(y_j/2s)^k}{k!} \frac{2^{-\theta_j/2-2-k} (z_j/s)^{\theta_j/2+k+1}}{\Gamma(\theta_j/2+2+k)} e^{-z_j/2s} \\
&= \frac{(z_i/2)^{\theta_i/2+1}}{\Gamma(\theta_i/2+1)} s^{-\theta_i/2-2-(n-1)} e^{-z_i/2s} \times \\
&e^{-(\vec{y}_i + \vec{z}_i)/2s} 2^{-\vec{\theta}_i/2-2-(n-1)} \prod_{j \neq i} \sum_{k=0}^{\infty} \frac{(y_j/2s)^k}{k!} \frac{2^{-k} (z_j/s)^{\theta_j/2+k+1}}{\Gamma(\theta_j/2+2+k)}.
\end{aligned}$$

We now exchange the product and the sum in the above. We will need some more notations for a compact representation. For any two vectors a and b , denote by

$$a^b = \prod_i a_i^{b_i}, \quad a! = \prod_i a_i!.$$

Also let $\Theta_i, \mathbf{y}_i, \mathbf{z}_i$ stand for the vectors $(\theta_j, j \neq i)$, $(y_j, j \neq i)$, and $(z_j, j \neq i)$ respectively.

Let $\mathbf{k}$ denote the vector $(k_j, j \neq i)$, where every k_j takes any nonnegative integer values. Let $\mathbf{k}'1$ be the sum of the coordinates of $\mathbf{k}$. Then

$$\begin{aligned}
&\prod_{j \neq i} \sum_{k=0}^{\infty} \frac{(y_j/2s)^k}{k!} \frac{2^{-k} (z_j/s)^{\theta_j/2+k+1}}{\Gamma(\theta_j/2+2+k)} \\
&= \sum_{N=0}^{\infty} (4s)^{-N} s^{-\vec{\theta}_i/2-N-(n-1)} \mathbf{z}_i^{\Theta_i/2+1} \sum_{\mathbf{k}'1=N} \frac{\mathbf{y}_i^{\mathbf{k}}}{\mathbf{k}!} \frac{\mathbf{z}_i^{\mathbf{k}}}{\prod_{j \neq i} \Gamma(\theta_j/2+2+k_j)}.
\end{aligned}$$

Thus, combining the expressions we get

$$\begin{aligned}
(22) \quad h_i(s) \prod_{j \neq i} q_s^{-\theta_j}(z_j, y_j) &= \frac{z_i^{\theta_i/2+1}}{\Gamma(\theta_i/2+1)} 2^{-\theta_i/2-1-\vec{\theta}_i/2-2-(n-1)} \times \\
&s^{-\theta_i/2-2-(n-1)} e^{-z_i/2s} e^{-(\vec{y}_i + \vec{z}_i)/2s} \sum_{N=0}^{\infty} 4^{-N} s^{-\vec{\theta}_i/2-2N-(n-1)} B_N,
\end{aligned}$$

where

$$B_N = \mathbf{z}_i^{\Theta_i/2+1} \sum_{\mathbf{k}'1=N} \frac{\mathbf{y}_i^{\mathbf{k}}}{\mathbf{k}!} \frac{\mathbf{z}_i^{\mathbf{k}}}{\prod_{j \neq i} \Gamma(\theta_j/2+2+k_j)}.$$

We can now integrate over s in (22) to obtain

$$\int_0^{\infty} h_i(s) \prod_{j \neq i} q_s^{-\theta_j}(z_j, y_j) ds = \sum_{N=0}^{\infty} B'_N \int_0^{\infty} s^{-a_N} e^{-b/s} ds,$$

where

$$(23) \quad B'_N = \frac{z_i^{\theta_i/2+1}}{\Gamma(\theta_i/2+1)} 2^{-\theta_0/2-2n+1} 4^{-N} B_N, \quad \theta_0 = \sum_{i=1}^n \theta_i,$$

$$(24) \quad a_N = \theta_i/2 + \vec{\theta}_i/2 + 2n + 2N = \theta_0/2 + 2n + 2N,$$

$$(25) \quad b = z_i/2 + (\vec{y}_i + \vec{z}_i)/2 = (\vec{y}_i + z_0)/2, \quad z_0 = \sum_{i=1}^n z_i.$$

Now a simple change of variable $w = 1/s$ shows

$$\begin{aligned} \int_0^\infty s^{-a_N} e^{-b/s} ds &= \int_0^\infty w^{a_N} e^{-bw} w^{-2} dw = \int_0^\infty w^{a_N-2} e^{-bw} dw \\ &= \frac{\Gamma(a_N-1)}{b^{a_N-1}} \int_0^\infty \frac{b^{a_N-1}}{\Gamma(a_N-1)} w^{a_N-2} e^{-bw} dw = \frac{\Gamma(a_N-1)}{b^{a_N-1}}. \end{aligned}$$

Since the i th coordinate of the exit point is zero, one can define $y_i = 0$ and $y_0 = \sum_{j=1}^n y_j = \vec{y}_i$ to simplify notation. Thus we obtain

$$\begin{aligned} \int_0^\infty h_i(s) \prod_{j \neq i} q_s^{-\theta_j}(z_j, y_j) ds &= \sum_{N=0}^\infty \frac{z_i^{\theta_i/2+1}}{\Gamma(\theta_i/2+1)} 2^{-\theta_0/2-2n+1} 4^{-N} B_N \frac{\Gamma(a_N-1)}{b^{a_N-1}} \\ &= \frac{z_i^{\theta_i/2+1} \mathbf{z}_i^{\Theta_i/2+1}}{\Gamma(\theta_i/2+1)} 2^{-\theta_0/2-2n+1} \sum_{N=0}^\infty ((\vec{y}_i + z_0)/2)^{-\theta_0/2-2n-2N+1} \\ &\quad \times \Gamma(\theta_0/2 + 2n + 2N - 1) 4^{-N} \sum_{\mathbf{k}'=1=N} \frac{\mathbf{y}_i^{\mathbf{k}}}{\mathbf{k}!} \frac{\mathbf{z}_i^{\mathbf{k}}}{\prod_{j \neq i} \Gamma(\theta_j/2 + 2 + k_j)} \\ &= \frac{\mathbf{z}^{\Theta/2+1}}{\Gamma(\theta_i/2+1)} 2^{-\theta_0/2-2n+1} \sum_{N=0}^\infty (y_0 + z_0)^{-\theta_0/2-2n-2N+1} 2^{\theta_0/2+2n+2N-1} \\ &\quad \times \Gamma(\theta_0/2 + 2n + 2N - 1) 4^{-N} \sum_{\mathbf{k}'=1=N} \frac{\mathbf{y}_i^{\mathbf{k}}}{\mathbf{k}!} \frac{\mathbf{z}_i^{\mathbf{k}}}{\prod_{j \neq i} \Gamma(\theta_j/2 + 2 + k_j)} \\ &= \frac{\mathbf{z}^{\Theta/2+1}}{\Gamma(\theta_i/2+1)} \sum_{N=0}^\infty (y_0 + z_0)^{-\theta_0/2-2n-2N+1} \times \\ &\quad \Gamma(\theta_0/2 + 2n + 2N - 1) \sum_{\mathbf{k}'=1=N} \frac{\mathbf{y}_i^{\mathbf{k}}}{\mathbf{k}!} \frac{\mathbf{z}_i^{\mathbf{k}}}{\prod_{j \neq i} \Gamma(\theta_j/2 + 2 + k_j)}. \end{aligned}$$

We have the following result.

Theorem 8. Let $Z_1, Z_2, \dots, Z_n$ be independent BESQ processes of dimensions $-\theta_1, \dots, -\theta_n$, where each $\theta_i \geq 0$. Assume that $Z_i(0) = z_i(0) > 0$, for every i .

The distribution of $(\tau, Z(\tau))$ is supported on the set $(0, \infty) \times \cup_{i=1}^n H_i$, where H_i is the subspace orthogonal to the i th canonical basis vector e_i . That is

$$H_i = \{(y_1, y_2, \dots, y_n) : y_i = 0\}.$$

- (i) Let G_i , $i = 1, 2, \dots, n$ be independent Gamma random variables with parameters $\theta_i/2 + 1$, $i = 1, 2, \dots, n$. The law of τ is the same as that of $\min_i \frac{z_i}{2G_i}$ and

$$P(\tau = T_i) = P\left(\frac{G_i}{z_i} > \frac{G_j}{z_j}, \text{ for all } j \neq i\right),$$

where T_i is the first hitting time of H_i .

- (ii) The restriction of the law of the random vector $Z(\tau)$, restricted to the hyperplane H_i , admits a density with respect to all the variables y_j 's, $j \neq i$, which is given by

$$(26) \quad \begin{aligned} &= \frac{S^{1-\theta_0/2-2n}}{\Gamma(\theta_0/2+1)} \prod_{j=1}^n z_j^{\theta_j/2+1} \sum_{N=0}^{\infty} \Gamma(\theta_0/2+2n+2N-1) S^{-2N} \\ &\times \sum_{\sum_{j \neq i} k_j = N} \prod_{j \neq i} \frac{(y_j z_j)^{k_j}}{k_j! \Gamma(\theta_j/2+2+k_j)}. \end{aligned}$$

Here

$$S = \sum_{i=1}^n (y_i + z_i), \quad y_i = 0, \quad \theta_0 = \sum_{i=1}^n \theta_i.$$

Using Theorem 4, we get that the exit distribution of $NWF(\delta_1, \dots, \delta_n)$ starting from a point $(z_1, \dots, z_n) \in \mathbb{S}_n$ is the image under the map

$$x_i \mapsto \frac{x_i}{\sum_{j=1}^n x_j}, \quad 1 \leq i \leq n$$

of the exit density of independent BESQ processes of dimensions $-\theta_1, \dots, -\theta_n$, where each $\theta_i = 2\delta_i$.

Theorem 9. *The exit density of $\mu \sim NWF(\delta_1, \dots, \delta_n)$ starting from $(z_1, \dots, z_n) \in \mathbb{S}_n$ is supported on the set $\cup_{i=1}^n F_i$, where F_i is the face $\{x \in \mathbb{S}_n : x_i = 0\}$, and admits the following description.*

- (i) Let G_i , $i = 1, 2, \dots, n$ be independent Gamma random variables with parameters $\delta_i + 1$, $i = 1, 2, \dots, n$. Then

$$(27) \quad P(\mu \text{ exits through } F_i) = P\left(\frac{G_i}{z_i} > \frac{G_j}{z_j}, \text{ for all } j \neq i\right),$$

- (ii) Let δ represent the vector $(\delta_1, \dots, \delta_n)$ and let $\delta_0 = \sum_{i=1}^n \delta_i$. The exit distribution of the process μ , restricted to F_i , admits a density with respect to all the variables x_j 's, $j \neq i$, which is given by

$$(28) \quad (\delta_i + 1) \sum_{N=0}^{\infty} \frac{\Gamma(N+n+\delta_0)}{\Gamma(N+2n+\delta_0)} \sum_{\sum_{j \neq i} k_j = N} \text{Dir}_n(z; \mathbf{k} + \delta + \mathbf{2}) \text{Dir}_{n-1}(x; \mathbf{k} + \mathbf{1}).$$

Here the inner sum above is over all nonnegative integers $(k_j, j \neq i)$ such that $\sum_{j \neq i} k_j = N$. The vector $\mathbf{k}$ represents a vector whose j th coordinate is k_j for all $j \neq i$, and $\mathbf{k}_i = 0$. The vectors $\mathbf{k} + \delta + \mathbf{2}$ and $\mathbf{k} + \mathbf{1}$ represent vector additions of $\mathbf{k}$, δ , and the vector of all twos, and $\mathbf{k}$ and the vector of all ones respectively. The factor Dir_{n-1} is a density with respect to the $(n-1)$ -dimensional vector $(x_j, j \neq i)$ with corresponding parameters $(\mathbf{k}_j + 1, j \neq i)$. It can also be interpreted as the conditional density of the n -dimensional $\text{Dir}_n(x; \mathbf{k} + \mathbf{1})$, conditioned on $x_i = 0$.

Note that the density in (28) is a mixture of Dirichlet densities, strikingly similar to those appearing as transition probabilities of the Wright-Fisher diffusions themselves. See Griffiths [8], Barbour, Ethier, and Griffiths [4], and Pal [12].

Proof of Theorem 9. This is a straightforward integration. We have assumed that $\sum_i z_i = 1$. Thus, $S = 1 + \sum_j y_j$; define $y_0 = \sum_j y_j$, and

$$x_j = y_j/y_0, \quad 1 \leq j \leq n.$$

Hence (26) simplifies to

$$(29) \quad \begin{aligned} &= \frac{(1+y_0)^{1-\theta_0/2-2n}}{\Gamma(\theta_0/2+1)} \prod_{j=1}^n z_j^{\theta_j/2+1} \sum_{N=0}^{\infty} \Gamma(\theta_0/2+2n+2N-1)(1+y_0)^{-2N} \\ &\times y_0^N \sum_{\sum_{j \neq i} k_j = N} \prod_{j \neq i} \frac{(x_j z_j)^{k_j}}{k_j! \Gamma(\theta_j/2+2+k_j)}. \end{aligned}$$

Now, to get to formula (28) we need to make a multivariate change of variables. Without loss of generality, let $i = n$. Then, for any $y \in F_i$, we have $y_n = 0$. Define the change of variables

$$(y_1, \dots, y_{n-2}, y_{n-1}) \mapsto (y_0, x_1, \dots, x_{n-2}).$$

In other words, $y_i = y_0 x_i$, for all $i = 1, 2, \dots, n-2$ and $y_{n-1} = y_0(1-x_1-\dots-x_{n-2})$. The determinant of the well-known Jacobian matrix is given by y_0^{n-2} .

Thus, the density of $(x_1, \dots, x_n)$ restricted to F_i is given by

$$(30) \quad \begin{aligned} &\frac{1}{\Gamma(\theta_i/2+1)} \prod_{j=1}^n z_j^{\theta_j/2+1} \sum_{N=0}^{\infty} \Gamma(\theta_0/2+2n+2N-1) \\ &\int_0^{\infty} y^{N+n-2} (1+y)^{1-\theta_0/2-2n-2N} dy \times \sum_{\sum_{j \neq i} k_j = N} \prod_{j \neq i} \frac{(x_j z_j)^{k_j}}{k_j! \Gamma(\theta_j/2+2+k_j)}. \end{aligned}$$

The following formula is easily verifiable for $\alpha \geq 0$, $\beta > \alpha + 1$:

$$\int_0^{\infty} y^{\alpha} (1+y)^{-\beta} dy = \int_0^1 x^{\beta-\alpha-2} (1-x)^{\alpha} dx = B(\alpha+1, \beta-\alpha-1)$$

where B refers to the Beta function.

In other words, (30) reduces to

$$(31) \quad \begin{aligned} &\frac{1}{\Gamma(\theta_i/2+1)} \prod_{j=1}^n z_j^{\theta_j/2+1} \sum_{N=0}^{\infty} \Gamma(\theta_0/2+2n+2N-1) \\ &B(N+n-1, N+n+\theta_0/2) \sum_{\sum_{j \neq i} k_j = N} \prod_{j \neq i} \frac{(x_j z_j)^{k_j}}{k_j! \Gamma(\theta_j/2+2+k_j)}. \end{aligned}$$

We now change $\theta_i/2$ to δ_i and rewrite the above expression in terms of Dirichlet densities. We use the notations in the statement of Theorem 9: the vector $\mathbf{k}$ represents a vector whose j th coordinate is k_j for all $j \neq i$, and $\mathbf{k}_i = 0$. The vectors $\mathbf{k} + \delta + \mathbf{2}$ and $\mathbf{k} + \mathbf{1}$ represent vector additions of $\mathbf{k}$, δ , and the vector of all twos, and $\mathbf{k}$ and the vector of all ones respectively. The factor Dir_{n-1} is a density with respect to the $(n-1)$ -dimensional vector $(x_j, j \neq i)$ with corresponding parameters $(\mathbf{k}_j + 1, j \neq i)$. It can also be interpreted as the conditional density of the n -dimensional $\text{Dir}_n(x; \mathbf{k} + \mathbf{1})$, conditioned on $x_i = 0$.

Hence, for any $(k_j, j \neq i)$, integers,

$$\begin{aligned} \frac{z_i^{\delta_i+1}}{\Gamma(\delta_i+1)} \prod_{j \neq i} \frac{z_j^{k_j+\delta_j+1}}{\Gamma(\delta_j+2+k_j)} \frac{x_j^{k_j}}{k_j!} = \\ \frac{(\delta_i+1)}{\Gamma(\delta_0+N+2n)\Gamma(N+n-1)} \text{Dir}_n(z; \mathbf{k} + \delta + \mathbf{2}) \text{Dir}_{n-1}(x; \mathbf{k} + \mathbf{1}). \end{aligned}$$

Thus (31) reduces to

$$(32) \quad \begin{aligned} (\delta_i+1) \sum_{N=0}^{\infty} \frac{\Gamma(\delta_0+2n+2N-1)B(N+n-1, N+n+\delta_0)}{\Gamma(\delta_0+N+2n)\Gamma(N+n-1)} \\ \times \sum_{\mathbf{k}' \mathbf{1}=N} \text{Dir}_n(z; \mathbf{k} + \delta + \mathbf{2}) \text{Dir}_{n-1}(x; \mathbf{k} + \mathbf{1}). \end{aligned}$$

However

$$\begin{aligned} & \frac{\Gamma(\delta_0+2n+2N-1)B(N+n-1, N+n+\delta_0)}{\Gamma(\delta_0+N+2n)\Gamma(N+n-1)} \\ &= \frac{\Gamma(\delta_0+2n+2N-1)}{\Gamma(\delta_0+N+2n)\Gamma(N+n-1)} \frac{\Gamma(N+n-1)\Gamma(N+n+\delta_0)}{\Gamma(2N+2n+\delta_0-1)} \\ &= \frac{\Gamma(N+n+\delta_0)}{\Gamma(N+2n+\delta_0)}. \end{aligned}$$

This completes the proof of formula (28).

The probability in (27) is a direct consequence of Theorem 8 conclusion (i). $\square$

5. EXIT TIME

Let $X = (X_1, \dots, X_n)$ be distributed as NWF $(-\theta_1/2, \dots, -\theta_n/2)$ starting from a point $(x_1, \dots, x_n)$ in the unit simplex. Let σ_0 denote the stopping time

$$\sigma_0 = \inf \{t \geq 0 : X_i = 0 \text{ for some } i\}.$$

Our objective is to find estimates on the law of σ_0 .

We will simplify the situation by assuming that all $x_i = 1/n$ and all $\theta_i = \theta$. To this end we use the time-change relationship in Theorem 4. Let $Z = (Z_1, \dots, Z_n)$ be independent BESQ processes starting from $(z_1, \dots, z_n)$ as in the set-up of Theorem 4, where each z_i is now one. Then

$$(33) \quad \sigma_0 = 4 \int_0^\tau \frac{ds}{\zeta(s)}, \quad \zeta(s) = \sum_{i=1}^n Z_i(s).$$

By Theorem 8, the distribution of τ is the same as considering n iid Gamma $(\theta/2+1)$ random variables $G_1, \dots, G_n$, and defining

$$(34) \quad \tau = \frac{1}{2 \max_i G_i}.$$

Our first step will be to prove a concentration estimate of $\max_i G_i$.

Lemma 10. *Let $G_1, G_2, \dots, G_n$ be n iid Gamma random variables with parameter $r/2$, for some $r \geq 2$. Let χ be the random variable $\max_i G_i$. Then, as n tends to infinity,*

$$E\sqrt{\chi} = \Theta\left(\sqrt{\log n}\right).$$

Proof. First let $r \in \mathbb{N}$. Let $\{Z_1(i), \dots, Z_n(i), i = 1, 2, \dots, r\}$ be a collection of iid standard Normal random variables. Then $2G_j$ has the same law as $Z_j^2(1) + \dots + Z_j^2(r)$. Hence

$$E \max_j |Z|_j(1) \leq E \sqrt{2\chi} \leq \sqrt{r} E \max_{i,j} |Z|_j(i)$$

As n tends to infinity, the right hand side above converges to $\sqrt{2r \log(rn)}$ while the left hand side converges to $\sqrt{2 \log n}$. This completes the argument for $r \in \mathbb{N}$. For a general positive r , bound on both sides by $\lfloor r \rfloor$ and $\lfloor r \rfloor + 1$. $\square$

We also need a version of logarithmic Sobolev inequality for Gamma random variables which can be found in several articles including [5].

Lemma 11. [5, p. 2718] *Let μ^θ denote the product probability measure of n iid $\text{Gamma}(\theta)$ random variables. Then, for every f on $\mathbb{R}^n$ which is in C^1 (i.e., once continuously differentiable) one has*

$$(35) \quad \text{Ent}(f^2) \leq 4 \int \left(\sum_{i=1}^n x_i (\partial_i f(x))^2 \right) d\mu^\theta(x).$$

Here $\text{Ent}(\cdot)$ refers to the entropy defined by

$$\text{Ent}(f^2) = \int f^2 \log(f^2) d\mu^\theta - \left(\int f^2 d\mu^\theta \right) \log \left(\int f^2 d\mu^\theta \right).$$

And ∂_i refers to the partial derivative with respect to the i th coordinate.

Lemma 12. *Consider the set-up in Lemma 11. Let F be a function on the open positive quadrant (i.e., every $x_i > 0$) which is C^1 and satisfies*

$$(36) \quad \sum_{i=1}^n x_i (\partial_i F)^2 \leq F.$$

Then the following concentration estimate holds for any $r > 0$:

$$\mu^\theta \left(\sqrt{F} - E_\theta \sqrt{F} \geq r \right) \leq \exp(-r^2), \quad \mu^\theta \left(\sqrt{F} - E_\theta \sqrt{F} \leq -r \right) \leq \exp(-r^2)$$

where $E_\theta \sqrt{F} = \int \sqrt{F} d\mu^\theta$.

Proof. Condition (36) implies that $4 \sum_{i=1}^n x_i \left(\partial_i \sqrt{F} \right)^2 \leq 1$. Hence, from the classical Herbst argument (for example, the monograph by Ledoux [11]), with a gradient defined by the right side of (35) we get

$$\mu^\theta \left(\sqrt{F} - E_\theta \sqrt{F} > r \right) \leq \exp(-r^2).$$

Here $\mu^\theta(\sqrt{F})$ is the expectation of $\sqrt{F}$ under μ^θ . Repeating the argument with $-\sqrt{F}$ instead of $\sqrt{F}$ we get the result. $\square$

Theorem 13. *The random variable $\chi = \max_i G_i$, where G_i 's are iid $\text{Gamma}(\theta)$, satisfies the following concentration estimate*

$$(37) \quad P(\sqrt{\chi} > E(\sqrt{\chi}) + r) \leq e^{-r^2}, \quad \text{for all } r > 0.$$

Proof of Theorem 13. To prove (37) we start by noting that Lemma 12 is satisfied by the family of $\mathbb{L}^k$ -norms, $\{F_k, k > 1\}$, defined by

$$F_k(x) = \left(\sum_{i=1}^n x_i^k \right)^{1/k}.$$

This is because each F_k is smooth (when every x_i is positive) and

$$(38) \quad \sum_{i=1}^n x_i (\partial_i F_k(x))^2 = \sum_{i=1}^n x_i \left[\frac{x_i^{k-1}}{\left(\sum_{j=1}^n x_j^k \right)^{1-1/k}} \right]^2 = \frac{\sum_{i=1}^n x_i^{2k-1}}{\left(\sum_{j=1}^n x_j^k \right)^{2-2/k}}.$$

Since for any nonnegative $y_1, y_2, \dots, y_n$ and any $\beta > 1$ one has

$$\sum_{i=1}^n y_i^\beta \leq \left(\sum_{i=1}^n y_i \right)^\beta,$$

applying it for $y_i = x_i^k$ and $\beta = 2 - 1/k$ we get

$$\sum_{i=1}^n x_i^{2k-1} \leq \left(\sum_{i=1}^n x_i^k \right)^{2-1/k}.$$

Combining the above with (38) we get

$$\sum_{i=1}^n x_i (\partial_i F_k(x))^2 \leq \left(\sum_{i=1}^n x_i^k \right)^{1/k} = F_k(x).$$

Thus F_k satisfies condition (36).

Since F_k converges pointwise to $\max_i x_i$ as k tends to infinity, by applying DCT, Lemma 12 is true for the function $\max_i G_i$. This proves (37). $\square$

Our next step will be to prove estimate on the quantity σ_0 in (33). The process $\zeta(s)$ is non-Markovian and not distributed as $Q^{-n\theta}$. However, on an possibly enlarged sample space one can create a $Q^{-n\theta}$ process $\tilde{\zeta}$ such that the paths of ζ and $\tilde{\zeta}$ are indistinguishable until σ_0 . This is possible by considering the SDE solved by ζ :

$$\zeta(t) = n - n\theta t + \int_0^t \sqrt{\zeta(s)} dW(s), \quad t < \sigma_0.$$

To extend the process beyond σ_0 , one concatenates an independent Brownian motion $\widetilde{W}$ and defines

$$\beta(t) = \begin{cases} W(t), & t \leq \sigma_0 \\ W(t) + \widetilde{W}(t - \sigma_0), & t > \sigma_0. \end{cases}$$

Then β is a Brownian motion in the enlarged filtration. Since $Q^{-n\theta}$ admits a strong solution, the process

$$(39) \quad \tilde{\zeta}(t) = n - n\theta t + 2 \int_0^t \sqrt{\tilde{\zeta}(s)} d\widetilde{W}(s), \quad t < T_0,$$

has law $Q^{-n\theta}$ and pathwise indistinguishable from ζ until time σ_0 . Thus in the following discussion we will treat as if ζ itself is distributed as $Q^{-n\theta}$ keeping in mind the above construction.

Theorem 14. *Let μ be distributed as an n -dimensional NWF($\delta, \delta, \dots, \delta$) starting from the point $(1/n, 1/n, \dots, 1/n)$. Let σ_0 be the first time that any of the coordinates of μ hit zero. Let*

$$a_n = E \max_{1 \leq i \leq n} \sqrt{G_i}, \quad G_i \stackrel{iid}{\sim} \text{Gamma}(\delta + 1).$$

Then, $a_n = \Theta(\sqrt{\log n})$, σ_0 has the law given by (33) where ζ is distributed as $Q_1^{-2n\delta}$ and τ is a random time.

Moreover, for any $r > 0$, we get

$$P\left(\frac{1}{n(a_n + r)} \leq \sqrt{2\tau} \leq \frac{1}{n(a_n + r)}\right) \geq 1 - 2e^{-r^2}.$$

Remark 1. It is impossible to provide a simple description of the exact distribution of σ_0 due to the distributional dependence of ζ and τ . The above theorem shows that τ is about a constant and one can compare the distribution of σ_0 with that of $4 \int_0^\cdot du/\zeta(u)$, where the upper limit of the integral is a constant. Limiting large deviation behavior of such integrals is possible to derive by methods as in [17].

Proof of Theorem 14. The proof is obvious from Lemma 13 and expression (34). $\square$

APPENDIX:

PROOFS OF PROPERTIES OF BESQ PROCESSES

Proof of Lemma 2. We use exercise 3.20 in [15, p. 311]. The scale function for Q^θ for $\theta \geq 0$ is well-known to be $x^{-\theta/2+1}$ (see [15, p. 443]). Nearly identical calculations lead to the case when θ is replaced by $-\theta$ and we obtain the scale function $s(x) = x^{\theta/2+1}$.

The speed measure is the measure with the density

$$m'(x) = \frac{2}{s'(x)4x} = \frac{1}{2(\theta/2 + 1)} x^{-\theta/2-1}.$$

We now Feller's criterion to check if the origin is an entry and/or exit point (see [9, p. 108]). Note that

$$(40) \quad m(\xi, 1/2) = \frac{1}{2(\theta/2 + 1)} \int_\xi^{1/2} x^{-\theta/2-1} dx = \frac{1}{\theta(\theta/2 + 1)} (\xi^{-\theta/2} - 2^{\theta/2})$$

$$m(0, \xi] = \infty, \quad \text{for all positive } \xi.$$

Thus

$$\int_0^{1/2} m(\xi, 1/2] s(d\xi) < \infty, \quad \text{and} \quad \int_0^{1/2} m(0, \xi] s(d\xi) = \infty.$$

This proves that the origin is an exit and not an entry.

Finally, to obtain part (ii) we apply Girsanov's theorem ([15, p. 327]). Let X satisfy the SDE $dX(t) = -\theta dt + 2\sqrt{X(t)}d\beta(t)$, then, we take $D(t) = X^{\theta/2+1}(t)$ (without the normalization, for simplicity) and apply Girsanov. Under the changed measure, there is a standard Brownian motion β^* such that

$$\begin{aligned} \beta(t) &= \beta^*(t) + \int_0^t X^{-\theta/2-1}(s) d\langle \beta, D \rangle_s \\ &= \beta^*(t) + \int_0^t X^{-\theta/2-1}(s) (\theta + 2) X^{\theta/2+1/2}(s) ds = \beta^*(t) + (\theta + 2) \int_0^t X^{-1/2}(s) ds. \end{aligned}$$

Thus, under the changed measure

$$dX(t) = -\theta dt + 2X^{1/2}(t)d\beta(t) = -\theta dt + 2(\theta + 2)dt + d\beta^*(t) = (\theta + 4)dt + d\beta^*(t).$$

The interpretation as the conditional distribution is classical (see [14]). $\square$

Proof of Lemma 3. For the assertion it is enough to take $t = 1$. Note that, under Q_0^θ , the coordinate process satisfies time-inversion, i.e., the process $\{t^2 Z(1/t), t \geq 0\}$ has law Q_0^θ . Thus, for $0 < \epsilon < 1$, if we define

$$U_\epsilon = \int_\epsilon^1 \frac{du}{Z(u)} = \int_1^{1/\epsilon} \frac{dt}{t^2 Z(1/t)},$$

then U_ϵ has the same law as $C_{1/\epsilon} - C_1 = \int_1^{1/\epsilon} du/Z(u)$. Thus, by [17, Thm. 1.1], we get $\lim_{\epsilon \rightarrow 0} U_\epsilon / \log(1/\epsilon) = (\theta - 2)^{-1}$ almost surely. $\square$

ACKNOWLEDGEMENT

I thank David Aldous, Zhen-Qing Chen, Michel Ledoux, and Jon Wellner for very useful discussions. I thank the anonymous referee for a thorough review which led to a significant improvement of the article.

REFERENCES

- [1] ALDOUS, D. (1993) The continuum random tree III. *The Annals of Probability* **21**, 248–289.
- [2] ALDOUS, D. (1999) Wright-Fisher diffusions with negative mutation rate! Available at <http://www.stat.berkeley.edu/~aldous/Research/OP/fw.html>.
- [3] ALDOUS, D. (2000) Mixing time for a Markov chain on cladograms. *Combin. Probab. Comput.* **9**, 191–204.
- [4] BARBOUR, A. D., ETHIER, S. N., and GRIFFITHS, R. C. (2000) A transition function expansion for a diffusion model with selection. *The Annals of Applied Probability* **10** (1), 123–162.
- [5] BARTHE, F. and WOLFF, P. (2009) Remarks on non-interacting conservative spin systems: The case of gamma distributions. *Stochastic Processes and their Applications* **119**, 2711–2723.
- [6] DURRETT, R. (2008) *Probability Models for DNA Sequence Evolution (Second Edition)*. Probability and its Applications. Springer.
- [7] GÖING-JAESCHKE and YOR, M. (2003) A survey and some generalizations of Bessel processes. *Bernoulli* **9** (2), 313–349.
- [8] GRIFFITHS, R. C. (1979) A transition density expansion for a multi-allele diffusion model. *Adv. Appl. Prob.*, **11** (2), 310–325.
- [9] ITÔ, K. and MCKEAN, H.P. (1996) *Diffusion processes and their sample paths*. Classics in Mathematics, Springer.
- [10] LAMBERT, A. (2010) The contour of splitting trees is a Lévy process. *The Annals of Probability* **38** (1), 348–395.
- [11] LEDOUX, M. (2000) *The Concentration of Measure Phenomenon*. Mathematical Surveys and Monographs **89**, American Mathematical Society.
- [12] PAL, S. (2009) Analysis of the market weights under the volatility-stabilized market models. *Preprint: accepted at The Annals of Applied Probability modulo minor revision*.
- [13] PAL, S. (2011) On the Aldous diffusion on continuum trees. I. Preprint available at [arXiv:1104.4186v1](https://arxiv.org/abs/1104.4186v1).
- [14] PITMAN, J. and YOR, M. (1982) A decomposition of Bessel bridges. *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **59**, 425–457.
- [15] REVUZ, D. and YOR, M. (1999) *Continuous Martingales and Brownian Motion, Third Edition* A Series of Comprehensive Studies in Mathematics **293**, Springer.
- [16] SCHWEINSBERG, J. (2000) An $O(n^2)$ bound for the relaxation time of a Markov chain on cladograms. *Random Structures and Algorithms* **20**(1), 59–70.
- [17] YOR, M. and ZANI, M. (2001) Large deviations for the Bessel clock. *Bernoulli* **7**(2), 351–362.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98115
E-mail address: `soumik@u.washington.edu`